## **V REVIEW OF STATISTICAL POWER**

The power of a statistical measure is defined as the probability of a significant observation given that an effect hypothesis  $(H_1)$  is true. Define the value of a dependent variable as X. Then, given that the null hypothesis  $(H_0)$  is true, a significant observation, x, is defined as one in which the probability of observing

$$x \ge \mu_0 + 1.645\sigma_0$$

where  $\mu_0$  and  $\sigma_0$  are the mean and standard deviation of the parent  $H_0$  distribution, is less than or equal to 0.05.

Figure 3 shows these definitions in graphical form under the assumption of normality. The *Z-Score* is a normalized representation of the dependent variable and is given by:

$$z = \frac{(x - \mu_0)}{\sigma_0},$$

where x is the value of the dependent variable and  $\mu_0$  and  $\sigma_0$  are the mean and standard deviation, respectively, of the parent distribution under  $H_0$ , and  $z_c$  is the minimum value (i.e., 1.645) required for significance (one-tailed). The mean of z under  $H_0$  is zero. The mean and standard deviation of z under  $H_1$  are  $\mu_{AC}$  and  $\sigma_{AC}$ , respectively.




Figure 3. Normal Representation of Statistical Power

This document is made available through the declassification efforts and research of John Greenewald, Jr., creator of:

## The Black Vault



The Black Vault is the largest online Freedom of Information Act (FOIA) document clearinghouse in the world. The research efforts here are responsible for the declassification of hundreds of thousands of pages released by the U.S. Government & Military.

**Discover the Truth at: http://www.theblackvault.com** 

## **Technical Protocol for the MEG Investigation**

Approved For Release 2003/09/09: CIA-RDP96-00789R003000260001-0

In general the effect size,  $\varepsilon$ , may be defined as:

$$\varepsilon = \frac{z}{\sqrt{n}},\tag{3}$$

where n is the sample size. Let  $\varepsilon_{AC}$  be the empirically derived effect size for anomalous cognition (AC). Then  $z_{AC} = \mu_{AC}$  in Figure 3 is computed from Equation 3. From Figure 3 we see that power is defined by:

Power = 
$$\frac{1}{\sigma_{AC}\sqrt{2\pi}} \int_{z_C}^{\infty} e^{-0.5 \left(\frac{\zeta - \mu_{AC}}{\sigma_{AC}}\right)^2} d\zeta.$$
 (4)

Let

$$z = \frac{\varsigma - \mu_{AC}}{\sigma_{AC}}.$$

Then Equation 4 becomes

Power = 
$$\frac{1}{\sqrt{2\pi}} \int_{z'c}^{\infty} e^{-0.5z^2} dz, \quad \text{where } z'_c = \frac{z_c - \mu_{AC}}{\sigma_{AC}}.$$
 (5)

For planning purposes, it is convenient to invert Equation 5 to determine the number of trials that are necessary to achieve a given power under the  $H_1$  hypothesis. If we define z(P) to be the z-score associated with a power, P, then the number of trials required is given by:

$$n = \frac{4z^2(P)}{\varepsilon_{AC}^2},\tag{6}$$

where  $\varepsilon_{AC}$  is the estimated mean value for the effect size under H<sub>1</sub>. Figure 4 shows the power, calculated from Equation 5, for various effect sizes for  $z_c = 1.645$ .




Figure 4. Statistical Power for Various Effect Sizes